

CAST search for sub-eV mass solar axions with ^3He buffer gas

S. Aune,¹ K. Barth,² A. Belov,³ S. Borghi,^{2,*} H. Bräuninger,⁴ G. Cantatore,⁵ J. M. Carmona,⁶ S. A. Cetin,⁷ J. I. Collar,⁸ T. Dafni,⁶ M. Davenport,² C. Eleftheriadis,⁹ N. Elias,² C. Ezer,⁷ G. Fanourakis,¹⁰ E. Ferrer-Ribas,¹ P. Friedrich,⁴ J. Galán,⁶ J. A. García,⁶ A. Gardikiotis,¹² E. N. Gazis,¹³ T. Gerasis,¹⁰ I. Giomataris,¹ S. Gninenko,³ H. Gómez,⁶ E. Gruber,¹¹ T. Guthörl,¹¹ R. Hartmann,^{14,†} F. Haug,² M. D. Hasinoff,¹⁵ D. H. H. Hoffmann,¹⁶ F. J. Iguaz,^{6,‡} I. G. Irastorza,⁶ J. Jacoby,¹⁷ K. Jakovčić,¹⁸ M. Karuza,⁵ K. Königsmann,¹¹ R. Kotthaus,¹⁹ M. Krčmar,¹⁸ M. Kuster,^{4,16,§} B. Lakić,^{18,¶} J. M. Laurent,² A. Liolios,⁹ A. Ljubičić,¹⁸ V. Lozza,⁵ G. Lutz,^{14,†} G. Luzón,⁶ J. Morales,^{6,**} T. Niinikoski,^{2,††} A. Nordt,^{4,16,‡‡} T. Papaevangelou,¹ M. J. Pivovarov,²⁰ G. Raffelt,¹⁹ T. Rashba,²¹ H. Riege,¹⁶ A. Rodríguez,⁶ M. Rosu,¹⁶ J. Ruz,^{6,2} I. Savvidis,⁹ P. S. Silva,² S. K. Solanki,²¹ L. Stewart,² A. Tomás,⁶ M. Tsagri,^{12,‡‡} K. van Bibber,^{20,§§} T. Vafeiadis,^{2,9,12} J. Villar,⁶ J. K. Vogel,^{11,20,¶¶} S. C. Yildiz,⁷ and K. Zioutas^{2,12}

(CAST Collaboration)

¹IRFU, Centre d'Études Nucléaires de Saclay (CEA-Saclay), Gif-sur-Yvette, France

²European Organization for Nuclear Research (CERN), Genève, Switzerland

³Institute for Nuclear Research (INR), Russian Academy of Sciences, Moscow, Russia

⁴Max-Planck-Institut für Extraterrestrische Physik, Garching, Germany

⁵Instituto Nazionale di Fisica Nucleare (INFN), Sezione di Trieste and Università di Trieste, Trieste, Italy

⁶Instituto de Física Nuclear y Altas Energías, Universidad de Zaragoza, Zaragoza, Spain

⁷Dogus University, Istanbul, Turkey

⁸Enrico Fermi Institute and KICP, University of Chicago, Chicago, IL, USA

⁹Aristotle University of Thessaloniki, Thessaloniki, Greece

¹⁰National Center for Scientific Research "Demokritos", Athens, Greece

¹¹Albert-Ludwigs-Universität Freiburg, Freiburg, Germany

¹²Physics Department, University of Patras, Patras, Greece

¹³National Technical University of Athens, Athens, Greece

¹⁴MPI Halbleiterlabor, München, Germany

¹⁵Department of Physics and Astronomy, University of British Columbia, Vancouver, Canada

¹⁶Technische Universität Darmstadt, IKP, Darmstadt, Germany

¹⁷Johann Wolfgang Goethe-Universität, Institut für Angewandte Physik, Frankfurt am Main, Germany

¹⁸Rudjer Bošković Institute, Zagreb, Croatia

¹⁹Max-Planck-Institut für Physik (Werner-Heisenberg-Institut), München, Germany

²⁰Lawrence Livermore National Laboratory, Livermore, CA, USA

²¹Max-Planck-Institut für Sonnensystemforschung, Katlenburg-Lindau, Germany

(Dated: June 21, 2011)

The CERN Axion Solar Telescope (CAST) has extended its search for solar axions by using ^3He as a buffer gas. At $T = 1.8$ K this allows for larger pressure settings and hence sensitivity to higher axion masses than our previous measurements with ^4He . With about 1 h of data taking at each of 252 different pressure settings we have scanned the axion mass range $0.39 \text{ eV} \lesssim m_a \lesssim 0.64 \text{ eV}$. From the absence of excess X-rays when the magnet was pointing to the Sun we set a typical upper limit on the axion-photon coupling of $g_{a\gamma} \lesssim 2.3 \times 10^{-10} \text{ GeV}^{-1}$ at 95% CL, the exact value depending on the pressure setting. KSVZ axions are excluded at the upper end of our mass range, the first time ever for any solar axion search. In future we will extend our search to $m_a \lesssim 1.15 \text{ eV}$, comfortably overlapping with cosmological hot dark matter bounds.

PACS numbers: 95.35.+d, 14.80.Mz, 07.85.Nc, 84.71.Ba

Introduction.—The Peccei-Quinn mechanism is the most compelling explanation for why in QCD the Θ term does not cause measurable CP-violating effects such as a large neutron electric dipole moment [1–3]. A testable consequence is the existence of axions, low-mass pseudoscalar bosons that are closely related to neutral pions. The axion mass is given by $m_a f_a \sim m_\pi f_\pi$ and the two-photon interaction strength scales with f_π/f_a where $f_\pi \sim 92 \text{ MeV}$ is the pion decay constant and f_a a large energy scale related to the breaking of a new U(1) sym-

metry of which the axion is the Nambu-Goldstone boson.

Axions would have been produced in the early universe by the vacuum realignment mechanism and radiation from cosmic strings, leading to a cold dark matter component, as well as from thermal interactions, leading to a hot dark matter component [4, 5]. Precision cosmology requires $m_a \lesssim 0.9 \text{ eV}$ for the latter [6, 7], with the usual caveats concerning systematic uncertainties. The cold component increases with decreasing m_a and provides all dark matter for $m_a \sim 10 \text{ } \mu\text{eV}$ ($f_a \sim 10^{12} \text{ GeV}$),




FIG. 1: Exclusion regions in the m_a - $g_{a\gamma}$ -plane achieved by CAST in the vacuum [20, 21], ^4He [22] and ^3He phase. We also show constraints from the Tokyo helioscope [17–19], horizontal branch (HB) stars [11], and the hot dark matter (HDM) bound [6]. The yellow band represents typical theoretical models with $|E/N - 1.95| = 0.07$ – 7 . The green solid line corresponds to $E/N = 0$ (KSVZ model).

with large uncertainties depending on the early-universe scenario. The ongoing ADMX dark matter search [8], based on Sikivie’s idea of axion-photon conversion in a macroscopic B field [9], provides one of the few realistic opportunities to find “invisible axions” [10].

Axions would also emerge from the hot interiors of stars, the Sun being the most powerful “local” source [11]. To search for these axions, one can use magnetically induced $a\gamma$ conversion in a dipole magnet pointing toward the Sun (“axion helioscope” technique [9]). This is analogous to neutrino flavor oscillations, $a\gamma$ mixing being caused by the B field [12]. The axion-photon interaction is given by Lagrangian $\mathcal{L}_{a\gamma} = g_{a\gamma} \mathbf{E} \cdot \mathbf{B} a$ with $g_{a\gamma} = (\alpha/2\pi f_a) [E/N - 2(4+z)/3(1+z)]$. Here $z = m_u/m_d$ with the canonical value 0.56, although the range 0.35–0.60 is possible [3]. E/N is a model-dependent ratio of small integers [13] and $E/N = 0$ (KSVZ model [14, 15]) is our benchmark case (green line in Fig. 1).

After a pioneering axion helioscope in Brookhaven [16], a fully steerable instrument was built in Tokyo [17–19]. The largest helioscope yet is the CERN Axion Solar Telescope (CAST), using a refurbished LHC test magnet ($L = 9.26$ m, $B \sim 9.0$ T) mounted to follow the Sun for about 1.5 h both at dawn and dusk [20–24]. CAST began operation in 2003 and after two years of data taking with vacuum inside the magnet bores achieved a limit of $g_{a\gamma} < 0.88 \times 10^{-10} \text{ GeV}^{-1}$ at 95% CL for $m_a \lesssim 0.02$ eV

[20, 21]. While these results are excellent to constrain very light axion-like particles [25], realistic QCD axions are not covered because the $g_{a\gamma}$ bounds quickly degrade for $m_a \gtrsim 0.02$ eV (Fig. 1).

Sensitivity to higher axion masses improves if the conversion volume contains a buffer gas such as helium [26]. Then the $a\gamma$ conversion probability is

$$P_{a \rightarrow \gamma} = \left(\frac{B g_{a\gamma}}{2} \right)^2 \frac{1 + e^{-\Gamma L} - 2e^{-\Gamma L/2} \cos(qL)}{q^2 + \Gamma^2/4} \quad (1)$$

where Γ is the inverse photon absorption length in the buffer gas, while the momentum difference between the a and γ propagation eigenstates is given by $q^2 = [(m_a^2 - m_\gamma^2)/2E]^2 + (g_{a\gamma} B)^2$. For $m_a = m_\gamma$, axions and photons are maximally mixed and reach $P_{a \rightarrow \gamma} = (g_{a\gamma} B L/2)^2 = 1.7 \times 10^{-17}$ for $L = 9.26$ m, $B = 9.0$ T and $g_{a\gamma} = 10^{-10} \text{ GeV}^{-1}$. For $m_a \neq m_\gamma$, the conversion probability rapidly decreases due to the axion-photon momentum mismatch.

The maximum $P_{a \rightarrow \gamma}$ can be restored by matching m_a with a photon refractive mass m_γ [26]. This method was first applied by the Brookhaven helioscope using ^4He as a buffer gas [16] and later allowed CAST to reach realistic axion models for $m_a \lesssim 0.4$ eV (Fig. 1) [22]. However, $T = 1.8$ K of the superconducting magnet restricts, due to condensation, the maximum ^4He pressure to ~ 14 mbar thus allowing us to scan axion masses $m_a \lesssim 0.4$ eV. To close the gap to the hot-dark matter bound, we have used ^3He as buffer gas to allow CAST to search up to $m_a \lesssim 1.15$ eV. The first results from this novel technique for the axion mass range $0.39 \lesssim m_a \lesssim 0.64$ eV are reported here.

Upgrades.— After completing the data taking with ^4He as a buffer gas, the CAST experiment performed several upgrades in order to prepare for data taking with ^3He . The most important upgrade was the design and installation of a sophisticated ^3He gas system.

To scan over a range of axion masses, CAST needs to control precisely the helium gas density in the cold bores. This is achieved by filling the cold bores with precisely metered amount of gas in incremental steps. The step size of the gas density is equivalent to a pressure change of between 0.083 mbar and 0.140 mbar (calculated for gas at nominal temperature of 1.8 K). To scan the whole available mass range efficiently, data taking runs cover two density settings per solar tracking. During the measurement, it is desired to have the gas density in the cold bores be as homogenous and stable as possible. To achieve the proper densities in the cold bores of CAST, and to be able to reproducibly refill the bores (allowing us to search the same axion mass), requires that the gas system is capable of adjusting to fluctuations of external conditions (e.g. variations of the room and magnet temperatures).

The ^3He system can be described as a hermetically closed gas circuit which is divided into functional sec-

tions with specific purposes: Storage, Trap purge system, Metering and ramping of gas density, Expansion volume, Recovery and circulation.

All the necessary helium for CAST physics runs is transferred to the storage volume that has been specifically engineered to keep the gas pressure below atmospheric. Before entering the metering volumes, the gas passes through two charcoal traps. The first one at ambient temperature traps oil and water vapour while the second at liquid nitrogen temperature removes residual gases.

The metering precision of the gas density is obtained by the accurate temperature control of the metering volumes, and by use of a metrology-grade pressure-measuring instruments to determine the amount of gas introduced into the cold bores. This amount of gas is calculated by accurately measuring the pressure decrease in the metering volumes. The reproducibility for the amount of gas sent from the metering volume into the magnet is 61 ppm.

The gas is confined in the cold bore region of the magnet with thin X-ray windows installed on both ends. The windows are made of 15 μm -thick polypropylene stretched over a mostly-open strongback structure to provide high X-ray transmission, resistance to a sudden rise in pressure and minimal helium leakage. Heaters on the window flanges allow for periodic bake-out of gases adsorbed on the polypropylene.

In case of quench, a sudden loss of superconductivity in the magnet, the temperature of the magnet increases rapidly. If the cold volume remains closed, the gas pressure abruptly increases and endangers the integrity of the X-ray windows. The windows can safely withstand pressures up to 1.2 bar, and to prevent rupture during a quench, the system must safely evacuate the ^3He from the cold bores to the expansion volume. Thus, the expansion volume, initially under vacuum, acts as a buffer reservoir for the gas that is intentionally expelled from the cold bores. The CAST ^3He system will be described in detail in a future publication.

It is a demanding task to compute the amount of gas needed to achieve the desired gas density. In fact, such calculations can only reliably be performed through computational fluid dynamic (CFD) simulations that account for the as-built system, as well as different physical phenomena such as hydrostatic effects, convection and buoyancy. The CFD simulations will be described in detail in a future publication.

During preparations for the ^3He data taking, the CAST X-ray detectors were upgraded as well. The Time Projection Chamber (TPC) with a multi-wire proportional readout [27] that had covered both bores of the sunset end of the magnet was replaced by two Micromegas detectors of similar dimensions of the one previously installed at the sunrise side [28] but with readouts fabricated with novel bulk and microbulk techniques [29–

31]. On the sunrise end a new shielded bulk (and later on microbulk) Micromegas replaced the unshielded one of our previous run [28]. These novel techniques provide several improvements in terms of stability and homogeneity of response, energy resolution, simplicity of construction [29–31] and, for the case of microbulk readouts, material radiopurity [32]. This is the first time these kinds of readouts are used in a physics run of a low background experiment. These new Micromegas detectors have obtained background levels down to $\sim 5 \times 10^{-6}$ counts $\text{keV}^{-1}\text{cm}^{-2}\text{s}^{-1}$ in the energy range of interest, one order of magnitude better than their predecessors [22]. This improvement is due to new shielding in the case of the sunrise detector, and to better rejection capabilities of the Micromegas readout with respect to the MWPC one, for the sunset set-up. The X-ray mirror telescope with a pn-CCD chip [33] covering the other bore of the sunrise side remained unchanged.

Data analysis and results.— Data presented in this paper correspond to the first 252 density steps of the ^3He phase, which encompass an equivalent axion mass range between 0.39 eV and 0.64 eV. The total available exposure time in axion-sensitive conditions is about 200 hours per detector, shared approximately equally among each of the four CAST detectors, as well as among the stated range of axion masses.

Data analysis is performed in a manner similar to our previous results obtained with ^4He gas. This time, however, we use an unbinned likelihood function that can be expressed as

$$\log L \propto -R_T + \sum_i^N \log R(t_i, E_i, d_i) \quad (2)$$

where the sum runs over each of the N detected counts and $R(t_i, E_i, d_i)$ is the event rate expected at the time t_i , energy E_i and detector d_i of the event i . R_T is the integrated expected number of counts over all exposure time, energy and detectors

$$R(t, E, d) = B_d + S(t, E, d) \quad (3)$$

where B_d is the background rate of detector d . $S(t, E, d)$ is the expected rate from axions in detector d which depends on the axion properties $g_{a\gamma}$ and m_a

$$S(t, E, d) = \frac{d\Phi_a}{dE} P_{a \rightarrow \gamma} \epsilon_d \quad (4)$$

where $P_{a \rightarrow \gamma}$ is the axion photon conversion probability in the CAST magnet (1), ϵ_d the detector efficiency, and

$$\frac{d\Phi_a}{dE} = 6.02 \times 10^{10} g_{10}^2 \frac{E^{2.481}}{e^{E/1.205}} \text{ cm}^{-2} \text{ s}^{-1} \text{ keV}^{-1} \quad (5)$$

is the solar axion spectrum, with $g_{10} = g_{a\gamma}/(10^{-10} \text{ GeV}^{-1})$ and energies in keV.

As explained in [22], the m_a dependency of the above expression is encoded in the probability $P_{a \rightarrow \gamma}$, which is coherently enhanced for values of m_a matching the photon mass m_γ induced by the buffer gas density, while it is negligible for values away from m_γ . Therefore, only the counts observed with the gas density matching a given axion mass m_a will contribute to the log L (and the exclusion plot) for that mass m_a .

The use of the unbinned likelihood (2), instead of the binned one used in our previous result [22] is motivated by the overall reduction of background rates achieved by CAST detectors with respect to the ones of the ^4He phase, as well as due to the reduced ^3He density setting exposure time (one half that for ^4He) due to time constraints of the overall data taking campaign. Indeed, the effective number of background counts in this analysis is about 1 count per density step for the Micromegas detectors, and about 0.2 in the fiducial spot of the CCD/Telescope system.

The remaining process is similar to the one followed in our previous results [22]: a best fit value $g_{\min}^4$ is obtained after maximization of L (for a fixed value of m_a). The obtained value is compatible with the absence of positive signal, and therefore an upper limit g_{95}^4 is obtained by integration of the Bayesian probability from zero up to 95% of its area in g^4 . This value is computed for many values of the axion mass m_a in order to configure the full exclusion plot shown in Fig. 1. A close up of the same exclusion plot is shown in Fig. 2, focused specifically in the axion mass range which has been explored in the data presented here.

As can be seen in Fig. 1, CAST extends its previous exclusion plot towards higher axion masses, excluding the interval 0.39–0.64 eV down to an average value of the axion-photon coupling of $2.27 \times 10^{-10} \text{ GeV}^{-1}$. The actual limit contour has high-frequency structure that is a result of statistical fluctuations that occur when a limit is computed for a specific mass using only a few hours of data.

Conclusions.—CAST has taken a great leap forward by using ^3He as buffer gas to cover m_a in the gap between our ^4He results and the hot dark matter bound. It is the first axion helioscope ever that has crossed the “axion line” for the benchmark KSVZ case. After covering $0.39 \text{ eV} \lesssim m_a \lesssim 0.64 \text{ eV}$ we will eventually reach 1.15 eV with the ^3He setup. If axions are not detected, the next challenge is to move down in the m_a – $g_{a\gamma}$ plot below the “axion band” of theoretical models. Such a goal cannot be achieved with the existing CAST apparatus and will require significant improvements of detector and magnet properties [34] or a completely new approach.

Acknowledgments.—We thank CERN for hosting the experiment and for the technical support to operate the magnet and cryogenics. We acknowledge support from NSERC (Canada), MSES (Croatia) under the grant number 098-0982887-2872, CEA (France), BMBF (Ger-




FIG. 2: Expanded view of the limit achieved in the ^3He CAST phase for axion mass range between 0.39 eV and 0.64 eV.

many) under the grant numbers 05 CC2EEA/9 and 05 CC1RD1/0 and DFG (Germany) under grant numbers HO 1400/7-1 and EXC-153, the Virtuelles Institut für Dunkle Materie und Neutrinos – VIDMAN (Germany), GSRT (Greece), RFFR (Russia), the Spanish Ministry of Science and Innovation (MICINN) under grants FPA2007-62833 and FPA2008-03456, NSF (USA) under Award number 0239812, US Department of Energy, NASA under the grant number NAG5-10842. Part of this work was performed under the auspices of the US Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344. We acknowledge the helpful discussions within the network on direct dark matter detection of the ILIAS integrating activity (Contract number: RII3-CT-2003-506222).

* Present addr.: Department of Physics and Astronomy, University of Glasgow, Glasgow, UK

† Present addr.: PNSensor GmbH, München, Germany

‡ Present addr.: IRFU, Centre d’Études Nucléaires de Saclay (CEA-Saclay), Gif-sur-Yvette, France

§ Present addr.: European XFEL GmbH, Notkestrasse 85, 22607 Hamburg, Germany

¶ Corresponding author: Biljana.Lakic@irb.hr

** Deceased.

†† Present addr.: Excellence Cluster Universe, Technische Universität München, Garching, Germany

‡‡ Present addr.: European Organization for Nuclear Research (CERN), Genève, Switzerland

§§ Present addr.: Naval Postgraduate School, Monterey, CA, USA

¶¶ Present addr.: Lawrence Livermore National Laboratory, Livermore, CA, USA

[1] R. D. Peccei, Lect. Notes Phys. **741**, 3 (2008).

[2] J. E. Kim and G. Carosi, Rev. Mod. Phys. **82**, 557 (2010).

[3] K. Nakamura *et al.* (Particle Data Group), J. Phys. G

- 37**, 075021 (2010).
- [4] P. Sikivie, Lect. Notes Phys. **741**, 19 (2008).
 - [5] O. Wantz and E. P. S. Shellard, Phys. Rev. D **82**, 123508 (2010).
 - [6] S. Hannestad, A. Mirizzi, G. G. Raffelt and Y. Y. Y. Wong, JCAP **1008**, 001 (2010).
 - [7] D. Cadamuro, S. Hannestad, G. Raffelt and J. Redondo, JCAP **1102**, 003 (2011).
 - [8] S. J. Asztalos *et al.* (ADMX Collaboration), Phys. Rev. Lett. **104**, 041301 (2010).
 - [9] P. Sikivie, Phys. Rev. Lett. **51**, 1415 (1983); (E) *ibid.* **52**, 695 (1984).
 - [10] S. J. Asztalos *et al.*, Ann. Rev. Nucl. Part. Sci. **56**, 293 (2006).
 - [11] G. G. Raffelt, Lect. Notes Phys. **741**, 51 (2008).
 - [12] G. Raffelt and L. Stodolsky, Phys. Rev. D **37**, 1237 (1988).
 - [13] S. L. Cheng, C. Q. Geng and W. T. Ni, Phys. Rev. D **52**, 3132 (1995).
 - [14] J. E. Kim, Phys. Rev. Lett. **43**, 103 (1979).
 - [15] M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Nucl. Phys. B **166**, 493 (1980).
 - [16] D. M. Lazarus *et al.*, Phys. Rev. Lett. **69**, 2333 (1992).
 - [17] S. Moriyama *et al.*, Phys. Lett. B **434**, 147 (1998).
 - [18] Y. Inoue *et al.*, Phys. Lett. B **536**, 18 (2002).
 - [19] Y. Inoue *et al.*, Phys. Lett. B **668**, 93 (2008).
 - [20] K. Zioutas *et al.* (CAST Collaboration), Phys. Rev. Lett. **94**, 121301 (2005).
 - [21] S. Andriamonje *et al.* (CAST Collaboration), JCAP **0704**, 010 (2007).
 - [22] E. Arik *et al.* (CAST Collaboration), JCAP **0902**, 008 (2009).
 - [23] S. Andriamonje *et al.* (CAST Collaboration), JCAP **0912**, 002 (2009).
 - [24] S. Andriamonje *et al.* (CAST Collaboration), JCAP **1003**, 032 (2010).
 - [25] J. Jaeckel and A. Ringwald, Ann. Rev. Nucl. Part. Sci. **60**, 405 (2010).
 - [26] K. van Bibber, P. M. McIntyre, D. E. Morris and G. G. Raffelt, Phys. Rev. D **39**, 2089 (1989).
 - [27] D. Autiero *et al.*, New J. Phys. **9**, 171 (2007).
 - [28] P. Abbon *et al.*, New J. Phys. **9**, 170 (2007).
 - [29] S. Andriamonje *et al.*, JINST **5**, P02001 (2010).
 - [30] J. Galan *et al.*, JINST **5**, P01009 (2010).
 - [31] S. Aune *et al.* (CAST Collaboration), Nucl. Instrum. Meth. A **604**, 15 (2009).
 - [32] S. Cebrian *et al.*, Astropart. Phys. **34**, 354 (2011).
 - [33] M. Kuster *et al.*, New J. Phys. **9**, 169 (2007).
 - [34] I. G. Irastorza *et al.*, JCAP **1106**, 013 (2011).